



## Sotatercept-csrk (WINREVAIR ™)

#### IMPORTANT REMINDER

We develop Medical Policies to provide guidance to Members and Providers. This Medical Policy relates only to the services or supplies described in it. The existence of a Medical Policy is not an authorization, certification, explanation of benefits or a contract for the service (or supply) that is referenced in the Medical Policy. For a determination of the benefits that a Member is entitled to receive under his or her health plan, the Member's health plan must be reviewed. If there is a conflict between the medical policy and a health plan or government program (e.g., TennCare), the express terms of the health plan or government program will govern.

#### **POLICY**

### **INDICATIONS**

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

#### FDA-Approved Indications

Winrevair is indicated for the treatment of adults with pulmonary arterial hypertension (PAH, World Health Organization [WHO] Group 1) to increase exercise capacity, improve WHO functional class (FC), and reduce the risk of clinical worsening events.

All other indications are considered experimental/investigational and not medically necessary.

### **DOCUMENTATION**

Submission of the following information is necessary to initiate the prior authorization review for initial requests: Chart notes, medical record documentation, or claims history supporting current pulmonary arterial hypertension (PAH) therapy.

#### PRESCRIBER SPECIALTIES

This medication must be prescribed by or in consultation with a pulmonologist or cardiologist.

### **COVERAGE CRITERIA**

#### **Pulmonary Arterial Hypertension (PAH)**

Authorization of 12 months may be granted for treatment of PAH in members 18 years of age and older when ALL of the following criteria are met:

- Member has PAH defined as WHO Group 1 class of pulmonary hypertension (refer to Appendix).
- PAH was confirmed by right heart catheterization with all of the following pretreatment (before any PAH therapy) results:
  - Mean pulmonary arterial pressure (mPAP) > 20 mmHg
  - Pulmonary capillary wedge pressure (PCWP) ≤ 15 mmHg
  - Pulmonary vascular resistance (PVR) >2 Wood units
- The requested medication will be used as add-on therapy.

This document has been classified as public information





- Member is currently receiving PAH therapy with medications from at least two of the following drug classes:
  - Endothelin receptor antagonist (e.g., Letairis, Opsumit, Tracleer)
  - Phosphodiesterase-5 inhibitor (e.g., Adcirca, Revatio)
  - Soluble guanylate cyclase stimulator (e.g., Adempas)
  - Prostacyclin analog (e.g., Flolan, Orenitram, Remodulin, Tyvaso, Veletri, Ventavis)
  - Prostacyclin receptor agonist (e.g., Uptravi)

#### **CONTINUATION OF THERAPY**

Authorization of 12 months may be granted for members with an indication listed in the coverage criteria section who are currently receiving the requested medication through a paid pharmacy or medical benefit, and who are experiencing benefit from therapy as evidenced by disease stability or disease improvement.

### **APPENDIX**

## WHO Classification of Pulmonary Hypertension (PH)

Note: Patients with heritable PAH or PAH associated with drugs and toxins might be long-term responders to calcium channel blockers.

### **Group 1: Pulmonary Arterial Hypertension (PAH)**

- Idiopathic
  - Long-term responders to calcium channel blockers
- Heritable
- Associated with drugs and toxins
- Associated with:
  - Connective tissue disease
  - Human immunodeficiency virus (HIV) infection
  - Portal hypertension
  - Congenital heart disease
  - Schistosomiasis
- PAH with features of venous/capillary(pulmonary veno-occlusive disease [PVOD]/pulmonary capillary hemangiomatosis [PCH]) involvement
- Persistent PH of the newborn

## **Group 2: PH associated with Left Heart Disease**

- Heart Failure:
  - With preserved ejection fraction
  - With reduced or mildly reduced ejection fraction
  - Cardiomyopathies with specific etiologies (i.e., hypertrophic, amyloid, Fabry disease, and Chagas disease)
- Valvular heart disease:
  - Aortic valve disease
  - Mitral valve disease
  - Mixed valvular disease
- Congenital/acquired cardiovascular conditions leading to post-capillary PH

## Group 3: PH associated with Lung Diseases and/or Hypoxia





- Chronic obstructive pulmonary disease (COPD) and/or emphysema
- Interstitial lung disease
- Combined pulmonary fibrosis and emphysema
- Other parenchymal lung diseases (i.e., parenchymal lung diseases not included in Group 5)
- Nonparenchymal restrictive diseases:
  - Hypoventilation syndromes
  - Pneumonectomy
- Hypoxia without lung disease (e.g., high altitude)
- Developmental lung diseases

## **Group 4: PH associated with Pulmonary Artery Obstructions**

- Chronic thromboembolic PH
- Other pulmonary artery obstructions
  - Sarcomas (high-or intermediate-grade) or angiosarcoma)
  - Other malignant tumors (e.g., renal carcinoma, uterine carcinoma, germ-cell tumors of the testis)
  - Non-malignant tumors (e.g., uterine leiomyoma)
  - Arteritis without connective tissue disease
  - Congenital pulmonary artery stenosis
  - Hydatidosis

## **Group 5: PH with Unclear and/or Multifactorial Mechanisms**

- Hematologic disorders, including inherited and acquired chronic hemolytic anemia and chronic myeloproliferative disorders
- Systemic disorders: Sarcoidosis, pulmonary Langerhans cell histiocytosis, and neurofibromatosis type 1
- Metabolic disorders, including glycogen storage diseases and Gaucher disease
- Chronic renal failure with or without hemodialysis
- Pulmonary tumor thrombotic microangiopathy
- Fibrosing mediastinitis
- Complex congenital heart disease

## **APPLICABLE TENNESSEE STATE MANDATE REQUIREMENTS**

BlueCross BlueShield of Tennessee's Medical Policy complies with Tennessee Code Annotated Section 56-7-2352 regarding coverage of off-label indications of Food and Drug Administration (FDA) approved drugs when the off-label use is recognized in one of the statutorily recognized standard reference compendia or in the published peer-reviewed medical literature.

#### **ADDITIONAL INFORMATION**

For appropriate chemotherapy regimens, dosage information, contraindications, precautions, warnings, and monitoring information, please refer to one of the standard reference compendia (e.g., the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) published by the National Comprehensive Cancer Network®, Drugdex Evaluations of Micromedex Solutions at Truven Health, or The American Hospital Formulary Service Drug Information).

### **REFERENCES**

- 1. Winrevair [package insert]. Rahway, NJ: Merck Sharp & Dohme LLC; March 2024.
- 2. Hoeper MM, Badesch DB, Ghofrani HA, et al. Phase 3 trial of sotatercept for treatment of pulmonary arterial hypertension. N Engl J Med. 2023;388(16):1478-1490. doi: 10.1056/NEJMoa2213558





- 3. Hoeper MM, Badesch DB, Ghofrani HA, et al. Phase 3 trial of sotatercept for treatment of pulmonary arterial hypertension. Supplementary appendix. N Engl J Med. 2023;Suppl Appendix.
- 4. Simonneau G, Montani D, Celermajer DS, et al. Hemodynamic definitions and updated clinical classification of pulmonary hypertension. Eur Respir J. 2019;53(1):1801913. doi:10.1183/13993003.01913-2018
- 5. Acceleron Pharma, Inc. A Study of Sotatercept for the Treatment of Pulmonary Arterial Hypertension (MK-7962-003/A011-11)(STELLAR). In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000- [4/25/2024]. Available from: https://clinicaltrials.gov/study/NCT04576988. NLM Identifier: NCT04576988.
- 6. Kovacs G, Bartolome S, Denton CP, et al. Definition, classification and diagnosis of pulmonary hypertension. Eur Respir J. 2024;64(4):2401324. doi: 10.1183/13993003.01324-2024
- 7. Chin KM, Gaine SP, Gerges C, et al. Treatment algorithm for pulmonary arterial hypertension. Eur Respir J. 2024;64(4):2401325. doi: 10.1183/13993003.01325-2024

**EFFECTIVE DATE** 5/31/2025

ID CHS